

Grading Udders

“Like many things, if you don’t measure it, you can’t and won’t manage it,” says Bob Weaber, University of Missouri Extension beef genetics specialist. “Udder quality represents one of the most important convenience traits in commercial beef production. Some would argue that udder quality represents an economic trait as poor udders require

additional labor inputs and thereby decrease profitability.

“As seedstock producers, you should manage udder quality closely,” Weaber adds. “Your commercial customers expect trouble-free cattle when they make a purchase at your farm or ranch. Don’t give your customers a reason to buy genetics from another source to ‘clean up’ udders produced by your bulls.”

He explains cows with good udder quality not only take less labor but also may have improved longevity due to lower incidence of mastitis or injury. “Poor udders, especially large teat size, may delay timely colostrum intake by calves, resulting in poorer transmission of passive immunity.”

For several years Hereford breeders have been asked

to report udder scores to the American Hereford Association (AHA) to be utilized in genetic evaluation.

In July the Beef Improvement Federation (BIF) adopted a two-score system for evaluating udders. See “Understanding the BIF Udder Scoring System.” The AHA Board of Directors followed suit and adopted the same system during its August meeting.

With the new system, producers score both teat size and udder suspension. The scores for both characteristics range from 1 to 9 (see next page).

“A two-score system allows independent evaluation of teat size and suspension characteristics,” Weaber says. “While there is a general trend for cows with large teat size to also have poorer suspension, this is not always the case. The independent scoring of the two important udder quality traits allows more precise measurement of each trait. Even though these scores are subjective, they’ll do a better job documenting variation in udder quality than a single-score system. When the two scores are used, the heritability estimates for each trait should be higher due to increased precision of measurement.”

Udder scores should be taken at calving – within the first 24 hours. “This is the time when poor udders will have the most swelling and edema,” Weaber says. “Calving is the most critical time for udder quality from a calf’s perspective – a 3-month-old calf may not have trouble nursing a ‘1’ teat size, but a 72-lb. newborn calf likely will.”

Weaber suggests producers get in the habit of scoring udders when they collect calving ease data. “Make udder score collection a routine part of your performance data collection,” he says. “Until you get comfortable with the scoring system, tape a scoring guide to your clipboard or reduce on a copy machine and tape in your ‘red book’ for use in the field. Try to get as close as you can to the guide and be as consistent as possible.

“Once you do it for a while, you’ll start picking out the really good ones and really bad ones easily. Be honest with yourself and use as much of the scale as needed. If she’s a ‘1’ on

What's the score?

Pictured below are nine udders. Try your hand at scoring the udders for suspension and teat size, and enter to win two \$50 gift certificates for Shop Hereford apparel. To participate, e-mail scores to adenton@hereford.org, fax to (785) 363-7203 or mail to the AHA headquarters, attention Udder Scoring Contest. Deadline for participation is Feb. 1. Winners and correct scores will be reported in the March issue.



1) Score = _____



2) Score = _____



3) Score = _____



4) Score = _____



5) Score = _____



6) Score = _____



7) Score = _____



8) Score = _____



9) Score = _____

Contestant name _____

Street address _____

City _____ State _____ Zip code _____

Phone number _____ e-mail address _____

Udder Scoring Contest

AMERICAN HEREFORD ASSOCIATION
P.O. BOX 21000
KANSAS CITY, MO 64111-0200
(816) 542-2727 FAX (816) 542-0821

REGISTRATION APPLICATION / BIRTH INFORMATION WORKSHEET

CALVIN

MEMBER # _____ BREEDER NAME _____

CHECK BOX FOR EACH ANIMAL YOU WANT TO TRANSFER ON ENTRY (complete information on back of form)

BREEDING INFORMATION				CALF INFORMATION														
DAM HERD ID OR REG NUMBER	SIRE HERD ID OR REG NUMBER	TYPE (BORN)	REG CALF (Y/N)	CALF HERD ID NUMBER	TATTOO		SEX	CALVING DATE			BIRTH WEIGHT	DAM UDDER SCORE	HAIR EZ	H/P/S	TWIN	BIRTH GROUP	VENTILATOR RECORD (HERD ID)	
					RIGHT EAR	LEFT EAR		MO	DAY	YR								
<input type="checkbox"/>	1																	

Hereford breeders record udder scores on the Registration Application/Birth Information Worksheet (Form 1).

suspension and '1' on teat size, give the scores. Your customers expect it!"

Submitting scores

Hereford breeders record udder scores on the Registration Application/Birth Information Worksheet (Form 1).

According to Jack Ward, AHA chief operating officer and director of breed improvement, with the new system, breeders will list the score in the same field used previously, but instead of recording one score, they will input the two digits. The first digit will be the udder suspension (attachment) score, and the second digit will be teat size. For example, if a cow has an attachment score of 7 and a teat size of 5, the breeder will record 75 in the "dam udder score" column.

If a cow has an attachment score of 7 and a teat size of 5, the breeder will record 75.

Ward also explains that the one-score udder score data previously collected by the Association is still valid and will be used in future evaluations. "We will be able to continue to use the old data with the new system because of the direct correlation of the two," he says.

Research shows udder and teat characteristics are heritable; thus, change can be made through selection.

"Consistent and steady progress can make dramatic improvements in udder quality," Weaber says. "Help yourself and your customers by selecting for and culling on udder quality. Improving and monitoring udder traits will pay dividends in the long run." **HW**



BIF FACTS

2008-10

Understanding the BIF Udder Scoring System

As the beef industry strives to become more competitive, producers are analyzing any and all inputs in an attempt to keep expenses under control. One aspect of this has been a more critical evaluation of the functional soundness of the cow herd. This can be likened to a "cow herd production quality audit." Why? Because a cow that isn't problem-free costs extra time and labor. Many refer to functional characteristics as convenience traits. Others call them necessities. It's a fact that most profitable operators are running their cows with less labor than ever before, making it even more essential that cows get it done without extra care.

Udder and teat quality are two of the most important functional traits. Udder and teat soundness are a concern for a number of reasons: 1. Labor associated with extra costs and reduced convenience; 2. Longevity, which may be reduced because of injury or mastitis; 3. Calf performance, affected by a reduction in milk flow, or lower colostrum intake by

* * *

Teat Size

Scores range from 9 (Very Small) to 1 (Very Large, balloon shaped) and are subjective assessments of teat length and circumference.

Oversized teats are difficult for newborn calves to nurse and the calf may not receive adequate colostrum, which could lead to a higher incidence of scours or decreased immunity levels in the newborn calf. In Canadian studies, cows with more desirable teat scores had calves that nursed sooner after birth.

Udder Suspension

Scores range from 9 (Very Tight) to 1 (Very Pendulous) and represent assessments of udder support.

Weak udder suspension results in pendulous udders that make it difficult for a calf to nurse. Weak suspension in the udder indicates a lack of support in the ligament that ties the udder to the cow's body wall. Over time, weakness in this ligament will allow the udder to hang down too far from the body and may subject the udder to serious problems and increased potential for injury.

How to Score

Udder and teat scores should be taken (on the weakest quarter) within 24 hours of calving. Scores should preferably be taken by the same person throughout the calving season.

newborn calves having difficulty nursing oversized teats; and 4. Udder and teat characteristics appear to be heritable. This means that there is definitely is variation in the udder quality of daughters from different sire groups. Thus, change can be made through selection.

Udder Suspension

Teat Size

Score



Very Tight



Very Small



Tight



Small



Intermediate/Moderate



Intermediate/Moderate



Pendulous



Large



Very Pendulous/Broken Floor



Very Large/Balloon Shaped

Photos courtesy of American Gelbvieh Association